

Content available at: <https://www.ipinnovative.com/open-access-journals>

International Journal of Clinical Biochemistry and Research

Journal homepage: <https://www.ijcbr.in/>

## Original Research Article

## Study of serum electrolytes, calcium and magnesium in preeclampsia at a tertiary hospital in central Karnataka

Basavaraj V Savadi<sup>1,\*</sup>, Archana H B<sup>2</sup>, Bhagyajyoti Nalwarkar<sup>3</sup>, Rashmi B M<sup>4</sup><sup>1</sup>Dept. of Biochemistry, Basaveshwara Medical College & Hospital, Chitradurga, Karnataka, India<sup>2</sup>Dept. of Biochemistry, D. Y. Patil Medical College and Hospital, Kolhapur, Maharashtra, India<sup>3</sup>Dept. of Obstetrics and Gynaecology, Vijayanagar Institute of Medical Sciences, Ballari, Karnataka, India<sup>4</sup>Dept. of Community Medicine, Basaveshwara Medical College & Hospital, Chitradurga, Karnataka, India

## ARTICLE INFO

## Article history:

Received 06-03-2021

Accepted 04-05-2021

Available online 23-07-2021

## Keywords:

Preeclampsia

Serum sodium

Potassium

Calcium

Magnesium

Central Karnataka

## ABSTRACT

**Background:** Preeclampsia(PE) is a hypertensive disorder affecting about 5-14% of pregnancies worldwide and around 20% of high-risk pregnancies, resulting in maternal and perinatal morbidity and mortality. Imbalances in milieu of serum sodium and potassium, that are involved in functioning of vascular smooth muscles, leads to vasoconstriction and hypertension in PE. Calcium deficiency stimulates parathyroid hormone, leading to vasoconstriction. Magnesium regulates blood pressure by monitoring vascular tone and structure. There is scarcity of studies on these factors in PE in central Karnataka.

**Aim:** To estimate levels of serum calcium, magnesium, sodium and potassium in preeclampsia.

**Settings and Design:** Hospital based cross-sectional study conducted from September, 2020 to February, 2021 at a Tertiary Teaching Hospital in Chitradurga.

**Materials and Methods:** 60 diagnosed cases of preeclampsia and equal number of age, parity and gestational age matched women with singleton normal pregnancies, attending antenatal clinic or admitted and treated in obstetrics wards of the hospital were included in the study. Clinical examination and laboratory investigations to estimate biochemical parameters were conducted.

**Statistical Analysis:** Data was analyzed in SPSS V:20. Statistical tests like chi-square and One-way ANOVA test were applied to find the significance of associations.

**Results:** Hypernatremia, hypokalemia, hypomagnesemia and hypocalcemia were significant in preeclampsia compared to normal pregnancy group. In severe preeclampsia, highest levels of serum sodium and lowest levels of serum potassium, magnesium and calcium were found compared to the mild PE group.

**Conclusion:** Timely estimation of serum electrolytes, calcium and magnesium through the course of pregnancy could help in early detection and prompt treatment of preeclampsia and also can prevent the adverse outcomes of pregnancies. Also, dietary restriction of sodium and dietary supplementation of calcium, magnesium and potassium rich foods to the pregnant women and calcium supplementation could reduce the risk of preeclampsia.

© This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## 1. Introduction

Preeclampsia is a common hypertensive disorder affecting about 5-14% of pregnancies worldwide and around 20% of high-risk pregnancies, resulting in maternal and perinatal morbidity and mortality.<sup>1</sup> Preeclampsia(PE) is

a triad of hypertension (blood pressure >140/90mmHg), oedema and proteinuria ( $\geq 300\text{mg}/24\text{h}$  or protein: creatinine ratio  $\geq 0.3$ ) including any one of: thrombocytopenia, elevated liver transaminases, pulmonary oedema, new onset renal insufficiency or cerebral or visual disturbances, occurring after 20 weeks of pregnancy.<sup>2</sup>

All through pregnancy, the metabolic demand of foetus increases by 2-3folds and various biochemical, metabolic

\* Corresponding author.

E-mail address: [bvsavdi69@gmail.com](mailto:bvsavdi69@gmail.com) (B. V. Savadi).

and haematological changes occur in pregnancy to support metabolism of both mother and growing foetus. Abnormal placentation in first trimester and maternal syndrome in second/third trimesters involving excess of antiangiogenic factors results in PE.<sup>3–11</sup>

Imbalances in milieu of serum sodium and potassium, that are involved in functioning of vascular smooth muscles, may lead to hyper-responsiveness of vasculature to vasoconstrictors like vasopressin, causing vasoconstriction and hypertension in PE.<sup>3–11</sup> Calcium deficiency stimulates parathyroid hormone and renin release and caused vasoconstriction. It also indirectly affects smooth muscle function by increasing magnesium levels. Magnesium regulates blood pressure by monitoring vascular tone and structure. Additionally, alteration in angiogenic and inflammatory markers also lead to altered vascular contraction and immune maladaptation. Thus, oxidative stress along with electrolyte and micronutrient imbalances are implicated in etiopathogenesis of PE.<sup>3,4,7–11</sup> As there is scarcity of studies on these above-mentioned factors in PE in central Karnataka, this study was undertaken at a tertiary level teaching hospital in Chitradurga, to estimate levels of serum calcium and magnesium, sodium and potassium in preeclampsia and to study the associations of these factors in preeclampsia compared to the normal singleton pregnancy group.

## 2. Material and Methods

After obtaining institutional ethics committee clearance, this hospital based comparative cross-sectional study was conducted from September 2020 to February 2021, at a tertiary level teaching hospital in Chitradurga. 60 diagnosed cases of preeclampsia and equal number of age, parity and gestational age matched women with singleton normal pregnancies, attending antenatal clinic or admitted and treated in obstetrics wards of the hospital during the study period were selected by simple random sampling method, and included after explaining the purpose of the study and obtaining their informed consent.

Preeclampsia was defined as blood pressure  $>140/90$ mmHg after 20 weeks of pregnancy and either proteinuria  $\geq 300$ mg/24h or protein:creatinine ratio  $\geq 0.3$  or one of the following: thrombocytopenia, elevated liver transaminases, pulmonary edema, new onset renal insufficiency, or cerebral or visual disturbances.<sup>2</sup>

Exclusion criteria consisted of pregnancy with renal disease, multiple pregnancy, gestational trophoblastic disease, pregnant women with chronic hypertension, diabetes, pregnancy with cardiovascular diseases, patients with history of acute or chronic liver diseases, patients taking corticosteroids, patients having chronic inflammatory disorders like rheumatoid arthritis, tuberculosis, osteoarthritis, inflammatory bowel disease, patients on drugs like diuretics, patients on magnesium or other metal

containing medications.

Information was collected by interview technique, clinical examination and investigations, and documented in a pre-formed semi-structured questionnaire that included participant's socio-demographic details, clinical history, natal history, anthropometric measurements of height and weight as per WHO guidelines.<sup>12</sup> Blood pressure was recorded in supine posture after a rest period of 10 minutes with the standard mercury sphygmomanometer, by a single observer. Average of two readings was noted.<sup>12</sup>

Under aseptic precautions 5 ml of venous blood sample was collected from antecubital vein in plain vacutainer both from study groups. This blood sample was allowed to clot and serum was separated by centrifugation at 5000rpm for 5 minutes. The serum was used to estimate sodium, potassium, magnesium and calcium levels. Biochemical analysis was done in the central laboratory of the hospital by in VITROS 4600 - Dry Chemistry Analyzer. Serum calcium was analyzed by arsenazo III method. Serum magnesium was analysed by dye binding method. Serum sodium and potassium were analysed by direct ISE method.

Preeclampsia patients were further classified as mild and severe preeclampsia cases as per the 2013 diagnostic criteria for preeclampsia by the task force of American College of Obstetrics and Gynecologists.<sup>13</sup>

### 2.1. Statistical analysis

Data was compiled in MS Office Excel spreadsheet and analysed using SPSS V: 20(Statistical Package for Social Service). Qualitative data variables were expressed in frequency and percentages. Quantitative data variables expressed in mean and standard deviation. Pearson's Chi-square test and one way ANOVA test were applied to find the significance of association between categorical and continuous variables of the 2 study groups respectively. Associations with p-value of  $<0.05$  were considered be statistically significant.

## 3. Results

The present study included 60 preeclamptic patients in preeclamptic group and 60 women with singleton normal pregnancies (comparative group). Among the preeclamptic patients, 75% were suffering from mild preeclampsia and 25% had severe preeclampsia. The average age of mild PE group was  $22.5 \pm 3.4$  years, average age of severe PE group was  $21.9 \pm 2.2$  and that of comparative group was  $23.8 \pm 2.6$  years. The study groups were comparable with respect to age, parity and gestational age ( $p > 0.05$ ). Pre-eclampsia was higher among primigravida (65% primigravida in mild PE and 73% primigravida in severe PE) when compared to multigravida women (35% in mild and 27% in severe PE). (Table 1)

**Table 1:** Characteristics of the study participants

Parameters	Normal singleton pregnancy group (n=60)	Pre-eclamptic group Mild = 45	Pre-eclamptic group Severe = 15	p-value
Age in years (mean ± SD)	23.8±2.6	22.5±3.4	21.9±2.2	0.712
<b>Gravida</b>				
Primigravida n (%)	59 (59%)	29 (65%)	11 (73%)	0.241
Multigravida n (%)	41 (41%)	16 (35%)	4 (27%)	
<b>Gestation age in weeks (mean ± SD)</b>	31.6 ± 3.8	32.4 ± 2.7	31.1±2.4	0.941

**Table 2:** Association of serum calcium, magnesium, sodium and potassium levels among preeclampsia and normal pregnancy groups

Parameters	Normal singleton pregnancy group (n=60)	Mild preeclampsia group (n=35)	Severe preeclampsia Group (n=15)	p-value
SBP in mm Hg (Mean±SD)	123.2 ± 9.7	158.2±11.6	161±5.4	<0.01
DBP in mmHg (Mean±SD)	81.4±2.6	96.6±4.9	114.5±4.7	<0.01
Serum Sodium (mmol/l) Mean±SD	139.2±5.4	142.4±6.2	147±3.4	<0.01
Serum Potassium (mmol/l) Mean±SD	4.1±0.6	3.8±0.7	3.5±0.4	<0.01
Serum Calcium (mg/dl) Mean±SD	9.7±0.7	8.3±0.7	7.9 ±0.4	<0.01
Serum Magnesium (mg/dl) Mean±SD	1.8±0.9	1.5±0.5	1.4±0.3	<0.01

(Note: Data are shown as Mean±SD; one way ANOVA test was applied for SBP, DBP, serum calcium and magnesium and Kruskal Wallis H test was applied for serum sodium and potassium)

**Table 3:** Risk factors of preeclampsia

Independent variables	B	S.E.	Sig.	Exp(B)	95% C.I.for EXP(B) :
Age in years	0.05	0.28	0.858	1.051	0.608-1.818
Gestational age in weeks	-0.031	0.188	0.868	0.969	0.67-1.402
BMI in kg/m <sup>2</sup>	-1.271	0.714	0.075	0.28	0.069-1.136
S.Calcium in mg/dl	-3.808	0.861	0.001	0.022	0.004-0.12
S.Magnesium in mg/dl	-5.029	2.322	0.03	0.007	0-0.62
S.Sodium in mmol/l	0.03	0.061	0.623	1.031	0.914-1.163
S.Potassium in mmol/l	-0.851	0.515	0.041	0.427	0.156-1.172
Constant	70.244	24.125	0.004	3.21E+30	


\*Multivariate logistic regression test applied (Dependent variable: Preeclampsia)

Table 2 shows the association of blood pressure and electrolyte levels among the 2 study groups. The average levels of systolic and diastolic blood pressure in both mild and severe preeclampsia group were significantly elevated compared to normal pregnancy group, with the highest average level of both SBP and DBP noted in the severe PE group SBP: 161±5.4mmHg and DBP: 114.5±4.7) (Table 2 )

Accordingly, the average serum sodium levels also showed an increasing trend, with the highest mean value noted in the severe preeclampsia group (147±3.4 mmol/l), followed by the mild preeclampsia group (142.4±6.2mmol/L), while normal pregnancy comparative group had the serum sodium levels in normal range (139.2±5.4mmol/l). These differences in the serum sodium

levels were found to be statistically significant.

Normal singleton pregnancy group had serum potassium, calcium and magnesium levels within the normal range (serum potassium: 4.1±0.6mmol/L, calcium: 9.7±0.7mg/dl, magnesium: 1.8±0.9mg/dl). These levels showed a declining trend among the mild (serum potassium: 3.8±0.7 mmol/L, calcium: 8.3±0.7mg/dl, magnesium: 1.5±0.5mg/dl) and severe PE groups, with lowest levels noted in the severe preeclampsia group (serum potassium: 3.5±0.4mmol/L, calcium: 7.9 ±0.4mg/dl, magnesium: 1.4±0.3mg/dL). These associations were found to be statistically significant. (Table 2 )



**Fig. 1:** Regression graph showing correlation of serum sodium and potassium, calcium and magnesium levels with mean arterial blood pressure among preeclampsia patients

#### 4. Discussion

The present study was conducted to find the association between serum electrolytes, serum calcium and magnesium among the mild and severe preeclamptic group compared to the normal singleton pregnancy group. In the study preeclampsia was more common among the primigravida (65% of mild preeclampsia and 73% of severe preeclampsia group were primigravida). Similar results are found in the study conducted by Guhan VN et al. Indumati V. et al.<sup>11,14</sup>

Alterations in serum electrolytes levels indicate the underlying physiological and pathological changes of pregnancy. In our study, hypernatremia and hypokalemia were found in both mild and severe preeclampsia with highest derangement seen in the severe preeclamptic group (highest sodium levels:  $147 \pm 3.4$  mmol/l and lowest potassium levels:  $3.5 \pm 0.4$  mmol/l). Similar results are found in the studies conducted elsewhere.<sup>6,9,11,15</sup> When pregnancy is complicated by pre-eclampsia there is an increased sensitivity of vasopressor substances like aldosterone and decreased production of cyclic GMP, endothelin and PGE2 and this may have implications in sodium retention and increased peripheral vascular resistance which further develop into hypertension. This increased sensitivity of aldosterone also affects potassium balance causing hypokalemia. Some studies of preeclampsia have shown abnormality in the transport of sodium and potassium

pump across the vascular smooth muscle cell membrane, which is normally responsible for maintenance of blood pressure.<sup>11,14</sup>

In the present study, hypomagnesemia was found among both the mild and severe preeclamptic patients (mild PE:  $1.5 \pm 0.5$  mg/dl.; severe PE:  $1.4 \pm 0.3$  mg/dL). Similar results are found in the studies conducted by Indumati V. et al and Sayyed AAK et al.<sup>11,14</sup> The success of magnesium therapy in the treatment for eclamptic seizures and its known effect on vascular responses in vitro suggest that magnesium is deficient in women with pre-eclampsia.<sup>16</sup> The possible mechanism of hypomagnesemia and pregnancy induced hypertension could be promotion of vascular muscle relaxation by magnesium.

In our study levels of mean serum calcium were significantly lower among both mild ( $8.3 \pm 0.7$  mg/dl) and severe preeclamptic group ( $7.9 \pm 0.4$  mg/dl) compared to normal pregnancy group ( $9.7 \pm 0.71$  mg/dl). The findings of our study are similar to studies conducted elsewhere.<sup>5,10,11,17</sup> Any imbalances in the serum calcium milieu as noted by increased calcium excretion are found to be associated with abnormalities in BP levels. Disrupted calcium homeostasis can lead to altered vasoconstrictions. Hypocalcaemia stimulates parathyroid hormone and renin release, which in-turn increase the intracellular calcium in smooth muscle cells, resulting in high sensitivity of

these cells to angiotensin II, further causing vasoconstriction and hypertension.<sup>5,10,11,17</sup> Further, there is an increase in demand for calcium with about 25grams of calcium required for foetal development, highlighting the importance of calcium in pregnancy.<sup>11,18</sup>

## 5. Conclusion

In preeclampsia water and electrolyte imbalances occur, leading to sodium retention and potassium depletion, that lead to peripheral vascular resistance and hypertension. Also, hypomagnesemia, hypocalcaemia are found to be significantly associated with preeclampsia. Hence, dietary restriction of sodium and dietary supplementation of calcium, magnesium and potassium rich foods to the pregnant women in the form of milk, cheese, soybean products, leafy vegetables during the course of pregnancy, could result in a reduction in the incidence of preeclampsia. Calcium supplementation could reduce the risk of preeclampsia as well as risk of preterm births. As magnesium has an essential role in both foetal and maternal wellbeing, magnesium also has to be supplemented in pregnancies. Thus, timely estimation of serum electrolytes, calcium and magnesium through the course of pregnancy could help in early detection and prompt treatment of preeclampsia and also can prevent the adverse outcomes of pregnancies.

## 6. Source of Funding

None.

## 7. Conflict of Interest

The authors declare that there is no conflict of interest.

## References

1. National Health Portal, Govt. of India, 2021. Available from: <https://www.nhp.gov.in/disease/gynaecology-and-obstetrics/preeclampsia>.
2. American College of Obstetricians and Gynecologists. Clinical management guidelines for obstetrician-gynecologists, ACOG practice bulletin number 202. Gestational hypertension and preeclampsia. *Obstet Gynecol.* 2019;133:1–25.
3. Rana S, Lemoine E, Granger JP, Karumanchi SA. Preeclampsia. *Circ Res.* 2019;124:1094–1112. doi:10.1161/circresaha.118.313276.
4. Khayat S, Fanaei H, Ghanbarzahi A. Minerals in Pregnancy and Lactation: A Review Article. *J Clin Diagn Res.* 2017;11(9):QE01–QE05. doi:10.7860/JCDR/2017/28485.10626.
5. Udenze IC, Arikawe AP, Azinge EC, Okusanya BO, Ebuehi OA. Calcium and Magnesium Metabolism in Pre-Eclampsia. *West Afr J Med.* 2014;33(3):178–82.
6. Prafula KM. Evaluation of serum electrolytes in preeclamptic pregnant women: A hospital based study. *Glob J Res Anal.* 2017;6:122–3.

7. Kanagal DV, Rajesh A, Rao K, Devi UH, Shetty H, Kumari S. Levels of serum calcium and magnesium in pre-eclamptic and normal pregnancy: A study from Coastal India. *J Clin Ciagn Res.* 2014;8(7):1.
8. Chhabra S, Singh A. Role of Calcium in Hypertensive Disorders of Pregnancy Current Status of Research a Mini Review. *J Nutr Disord Ther.* 2017;7(2):1–5.
9. Gupta M, Roy N. Serum Sodium and Potassium Levels in Preeclampsia: A Clinical Study. *Indian J Med Biochem.* 2018;22(2):105–7. doi:10.5005/jp-journals-10054-0065.
10. Mittal S, Shaikh M, Thakur R, Jain D. Comparison of serum calcium and magnesium levels between preeclamptic and normotensive healthy pregnant women. *Int J Reprod Contracept Obstet Gynecol.* 2014;3(4):959–62. doi:10.5455/2320-1770.ijrcog20141216.
11. Indumati V, Kodliwadmath MV, Sheela MK. The role of serum electrolytes in pregnancy induced hypertension. *J Clin Diagnostic Res.* 2011;5(1):66–75.
12. Riley L, Guthold R, Cowan M, Savin S, Bhatti L, Armstrong T, et al. The World Health Organization STEPwise Approach to Noncommunicable Disease Risk-Factor Surveillance: Methods, Challenges, and Opportunities. *Am J Public Health.* 2016;106:74–8. doi:10.2105/ajph.2015.302962.
13. American College of Obstetricians and Gynecologists, Task Force on Hypertension in Pregnancy:“Report of the American College of Obstetricians and Gynecologists’ Task Force on Hypertension in Pregnancy. *Obstet Gynecol.* 2013;p. 1122.
14. Sayyed AAK, Pratinidhi SA. High sensitive C reactive protein as an inflammatory indicator in preeclampsia. *Int J Res Med Sci.* 2020;8(8):3002. doi:10.18203/2320-6012.ijrms20203454.
15. Manjareeka MA, Nanda SI. Serum electrolyte levels in preeclamptic women: a comparative study. *Int J Pharma and BioSci.* 2012;3(2):572–8.
16. Idogun ES, Imarengiaye CO, Momoh SM. Extracellular Calcium and Magnesium in Preeclampsia and Eclampsia. *Afr J Reprod Health.* 2007;11(2):89. doi:10.2307/25549719.
17. Ritchie LD, King JC. Dietary calcium and pregnancy-induced hypertension: is there a relation? *Am J Clin Nutr.* 2000;71(5):1371S–4S. doi:10.1093/ajcn/71.5.1371s.
18. Crowther CA, Hiller JR, Pridmore B, Bryce R, Duggan P, Hague WM. Calcium Supplementation In Nulliparous Women For The Prevention Of Pregnancy-Induced Hypertension, Preeclampsia And Preterm Birth: An Australian Randomized Trial. *Aust New Zealand J Obstet Gynaecol.* 1999;39(1):12–8. doi:10.1111/j.1479-828x.1999.tb03434.x.

## Author biography

**Basavaraj V Savadi**, Associate Professor

**Archana H B**, Assistant Professor

**Bhagyajyoti Nalwarkar**, Postgraduate Student

**Rashmi B M**, Associate Professor

**Cite this article:** Savadi BV, Archana H B, Nalwarkar B, Rashmi B M. Study of serum electrolytes, calcium and magnesium in preeclampsia at a tertiary hospital in central Karnataka. *Int J Clin Biochem Res* 2021;8(2):145-149.