Content available at: <https://www.ipinnovative.com/open-access-journals>

International Journal of Clinical Biochemistry and Research

Journal homepage: <https://www.ijcbr.in/>

Original Research Article

Adaptogens as anti-stress agents in reducing increased plasma cortisol level during stress

Anju^{1,*}, Ashis Kumar Ghosh¹¹Dept. of Biochemistry, Patna University, Patna, Bihar, India

ARTICLE INFO

Article history:

Received 27-08-2021

Accepted 23-09-2021

Available online 08-10-2021

Keywords:

Adaptogens

Bacopa monnieri

Cortisol

Hypothalamic-pituitary-adrenal axis

Ocimum sanctum

Panax ginseng

Withania somnifera

ABSTRACT

The stress response involves the activation of both the sympathetic-adrenal response and the hypothalamic-pituitary-adrenal axis. During times of increased stress, the adrenal gland is stimulated to produce increased levels of hormones. Cortisol, the main hormone involved in the stress response, is secreted in increased amounts within minutes of a perceived stressor. Cortisol secretion can increase as much as 20-fold and has several important physiological effects. Short-term surges in cortisol levels can suppress inflammation and at the same time suppress immune function. Though inflammation control is important, surges of sustained levels of cortisol are not healthy and ultimately lead to premature aging, degenerative disease, and increased susceptibility to cancer. Studies show that psycho-social stress activates the hypothalamus-pituitary-adrenal axis causing an increase in morning cortisol levels, which correlated to the subjects reports of increased fatigue and anxiety. Although this stress response is important for survival during an acute stressor, prolonged activation of the stress response may lead to adrenal exhaustion in which cortisol levels drop to insufficient levels resulting in fatigue or illness. Many herbs have been shown to impact adrenal function. Adaptogens are plants that produce a non-specific response improving the physiological resistance to stressors. These herbs are often used in the context of adrenal support formulas to balance adrenal hormone levels. It is believed that adaptogenic herbs can increase low levels of adrenal hormone or decrease levels that are elevated. Additionally, these herbs provide balancing activity on many body systems that are impacted by stress, such as the immune response and blood sugar control.

In the present study, we have evaluated the efficacy of ethanolic extract of *Ocimum sanctum* 47mg/kg p.o, *Withania somnifera* 23 mg/kg p.o and *Bacopa monnieri* 23 mg/kg p.o on plasma cortisol level in mice subjected to swim endurance test and cold restraint stress. The standard group was administered water-soluble root powder of *Panax ginseng* 100 mg/kg p.o and the stress control group was administered distilled water orally for 7 days. It was found that mice pretreated with ethanolic extracts of *Ocimum sanctum*, *Withania somnifera* and *Bacopa monnieri* showed a fall in the plasma cortisol level. The standard group also showed a significant decrease in the plasma cortisol level compared to the stress and normal control groups.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Adaptogens are natural substances found only in a few rare plants and herbs that normalize body functions, strengthen systems and functions compromised by stress and have a

protective effect against a wide variety of environmental and emotional stress. Ultimately, they are proposed to reduce chronic stress and fatigue through the adaptation of stress. Adaptogens have been around in Chinese functional medicine and Indian Ayurveda medicine since ancient times.¹ They became more popular during world war II when Russian scientists were looking to improve soldiers

*Corresponding author.

E-mail address: anju.fr@gmail.com (Anju).

stamina through herbal medicine.²⁻⁹

Adaptogens are similar to catecholamines, neurotransmitters involved in stress situations. Common catecholamines are adrenaline, epinephrine, dopamine and norepinephrine. Since, adaptogens are mild stressors the theory is that they build the body's immunity up to be able to adapt and control future severe and moderate stress situations. These stress-protective effects mainly help the Hypothalamus-Pituitary-Adrenal Axis and SAS system. They are responsible for controlling the body's stress responses during times of stress – mental health disorders, traumatic injury, exercise, eating disorders, surgery, malnutrition and low blood sugar amongst other conditions.¹⁰ Adaptogenic drugs are found in the most diverse families, they are distinctly different in their pattern of constituents.

Adaptogens work naturally to help regulate proper levels of stress-induced hormones and maintain healthy, balanced metabolic function.¹¹

Holy basil (*Ocimum sanctum*) is an adaptogenic herb, one of the primary botanicals used in Ayurvedic medicine to modulate a physiological stress response and to increase adaptive energy. Traditional Ayurvedic use for anti-stress activity has been verified by numerous animal clinical studies.^{12,13} A variety of studies have shown sacred basil to support the health of the cardiovascular, immune and digestive systems acting as an effective remedy to help ease stress-related conditions, including hypertension, lowering cortisol levels, mood disorders and peptic ulcers.¹²⁻¹⁵

Ashwagandha (*Withania somnifera*) has antibacterial, antitumor, anti-inflammatory and immunomodulating properties. A separate study also showed immunomodulatory (harmonizes immune function) effects, including a positive effect on the central nervous system. These tests show strong anti-stress actions, increasing memory and learning capabilities.¹⁶ The active constituents of *Withania somnifera* have been shown to reduce or modulate increased levels of circulating adrenocorticotrophic hormone (ACTH) and the concurrent increase in glucocorticoid levels (cortisol) to normal levels which may protect the body from the deleterious effects of chronic stress.¹⁷⁻²⁰

Brahmi (*Bacopa monnieri*) is used in a vast range of diseases. Brahmi is one of the best nerve tonic and is widely used in treating epilepsy, hysteria etc. Brahmi acts as a powerful brain food, and is known for its ability to enhance mind power.^{21,22} It supports and improves all aspects of mental functioning including comprehension, memory and recollection.²¹⁻²⁴ It is important because it also enhances crucial coordination of these three aspects of mental functioning. As it increases the ability to solve problems effectively, Brahmi is often used in formulas to help relieve and prevent stress.²⁵⁻²⁸

2. Materials and Methods

Plant material leaves of *Ocimum sanctum*, *Withania somnifera* and roots of *Bacopa monnieri* were collected, dried in shade, and finely powdered. The powder was soaked in absolute ethanol (95%) and left for 48 hours. The supernatant was collected and the residue was further soaked in absolute ethanol (95%) for 24 hours. The supernatant was collected and filtered. The filtrate was subjected to Rotavapour extraction at a temperature below 60°C for 24 hours. The concentrated form of the extract was obtained and freeze-dried.

The study was conducted on healthy, adult, male albino mice having a body

weight of 35 ± 5 g. They were acclimatized to laboratory condition for 2 weeks prior to experimentation. Animals were housed in polypropylene cages (6 mice/cage) in a mice experimentation laboratory at a temperature of 25°C ± 2°C with 12 – 12 h dark - light cycle. They were provided with standard food and water ad libitum. Institutional animal ethical committee (I.A.E.C) approval was obtained before the experiment and care was taken to handle the mice in humane manner. All the chemicals used in the present study were obtained from Euro Diagnostics (Mumbai, India), India Scientific Company (Patna, Bihar) and Bihar Scientific Corporation (Patna, Bihar).

3. Experimental

The adult animals (8 weeks old) were divided into 4 groups (n = 6 in each group) as follows: Group I consisted of Normal control (NC), these mice remained undisturbed in the home cage throughout the experimental period. Group II consisted of Stress control (SC), which were fed with equivolume of distilled water orally for 7 days. Group III (Stress+P.ginseng) consisted the standard group, these mice were fed with aqueous root powder of *Panax ginseng* (p.o) for 7 days.

Group IV consisted of (Stress+O.sanctum), (Stress+W.somnifera), (Stress+B.monneri) treatment group which were fed with ethanolic extract of *Ocimum sanctum*, *Withania somnifera*, *Bacopa monnieri* (p.o) for 7 days.

3.1. Stress procedure

3.1.1. Swim endurance test

The mice in group IV were given ethanolic extract of *Ocimum sanctum* 47 mg/kg (p.o), for 7 days. The standard group (III) was administered water soluble root powder of *Panax ginseng* 100 mg/kg (p.o), while the stress control group (II) was administered distilled water orally, for 7 days.

On the 8th day, the animals were allowed to swim till exhausted in a polypropylene tank of dimension 24 cm* 17 cm* 14 cm, filled with water to a height of 10 cm. The end point was taken when the animals drowned and 'swimming

time' for each animal was noted. The mean swimming time for each group was calculated and the data was statistically analyzed (Kumar et al., 1999).

3.1.2. Cold restraint stress

The mice in group IV were given ethanolic extract of *Ocimum sanctum* 47 mg/kg, *Withania somnifera* 23 mg/kg, *Bacopa monnieri* 27 mg/kg (p.o), for 7 days. The standard group (III) was administered water soluble root powder of *Panax ginseng* 100 mg/kg (p.o), while the stress control group (II) was administered distilled water for 7 days, orally.

On the 8th day, the animals were individually placed in plastic containers of capacity 350 ml. They were immobilized in their normal position, using adhesive tape. The containers were placed in a cold chamber maintained at 4°C for 2 hours. The blood was collected by orbital sinus veinpuncture method in a heparinised tube and plasma cortisol was determined by Enzyme Linked Immunosorbent Assay (ELISA).

3.2. Statistical analysis

Data was analyzed by the application of One way analysis of variance (ANOVA) using Graph pad in stat software. $P < 0.01$ was considered to be significant.

4. Result

Acute toxicity studies with extract revealed that LD₅₀*Ocimum sanctum* is 4.5g /kg, LD₅₀*Withania somnifera* is 1750 mg/kg, LD₅₀*Bacopa monnieri* is 17g/kg body weight (p.o). As shown in Figure 1, the extract of *Ocimum sanctum* improves swim duration in mice. Mice pretreated with ethanolic extract of *Ocimum sanctum* 47 mg/kg *Withania somnifera* 23 mg/kg *Bacopa monnieri* 27 mg/kg and water soluble root powder of *Panax ginseng* 100mg/kg (p.o) show significant improvement in the swimming time ($P < 0.01$), as compared to control. (n = 6 in all groups, SC vs S+O.sanctum, $P < 0.01$; SC vs S+P.ginseng, $P < 0.01$; One way ANOVA, $P < 0.01$, $F = 41.336$). (n = 6 in all groups, SC vs S+W.somnifera, $P < 0.01$; SC vs S+P.ginseng, $P < 0.01$; One way ANOVA, $P < 0.01$, $F = 41.336$). (n = 6 in all groups, SC vs S+B.monniieri, $P < 0.01$; SC vs S+P.ginseng, $P < 0.01$; One way ANOVA, $P < 0.01$, $F = 41.336$).

The induction of cold restraint stress led to a rise in plasma cortisol level. All the four treatments produced a significant reduction in plasma cortisol level. The plasma cortisol level which was found to be elevated in the animals subjected to cold restraint stress was significantly reduced by all the four treatments ($P < 0.01$), compared to controls. (n = 6 in all groups, NC vs SC, $P < 0.01$; SC vs S+O.sanctum, $P < 0.01$; SC vs S+P.ginseng, $P < 0.01$; One way ANOVA, $P < 0.01$, $F = 92.616$). (n = 6 in all groups, NC vs SC, $P < 0.01$; SC vs S+W.somnifera, $P < 0.01$; SC vs S+P.ginseng, $P < 0.01$;

Fig. 1: Effect of ethanolic extract of *Ocimum Sanctum* and aqueous root powder of *panax ginseng* on swimming performance in mice

Fig. 2: Effect of ethanolic extract of *Withania somnifera* and aqueous root powder of *panax ginseng* on swimming performance in mice

One way ANOVA, $P < 0.01$, $F = 92.616$). (n = 6 in all groups, NC vs SC, $P < 0.01$; SC vs S+B.monniieri, $P < 0.01$; SC vs S+P.ginseng, $P < 0.01$; One way ANOVA, $P < 0.01$, $F = 92.616$).

5. Discussion

Though the hypothalamus is traditionally considered the source of corticotropin-releasing factor (CRF), this neurotransmitter hormone and its receptors are also found in the gastrointestinal tract, skin, adrenal gland, testis, cardiac muscle, thymus and spleen.^{11,20,29} These particular tissues are also involved in the development of disorders related to heightened stress sensitivity and dysregulation of stress-coping mechanisms such as anorexia, diabetes, loss of libido, heart disease and immune pathologies.²⁵ It is well-established that CRF is negatively regulated by cortisol and that heightened stress sensitivity and dysregulation of stress-coping mechanisms appear to involve regulatory mechanisms of CRF.³⁰

Fig. 3: Effect of ethanolic extract of *Bacopa monnieri* and aqueous root powder of *Panax ginseng* on swimming performance in mice

Fig. 4: Effect of ethanolic extract of *Ocimum sanctum* and aqueous root powder of *Panax ginseng* on cold restraint stress induced changes in Plasma cortisol level in mice

Fig. 5: Effect of ethanolic extract of *Withania Somnifera* aqueous root powder of *Panax ginseng* on cold restraint stress induced changes in plasma cortisol level in mice

Fig. 6: Effect of ethanolic extract of *Bacopa monnieri* and aqueous root powder of *Panax ginseng* on Cold restraint stress induced changes in plasma cortisol level in mice

It has been shown that some herbal formulas can inactivate the secretion of CRF in rats by inhibiting CRF mRNA (mitochondrial) expression. Preventing the synthesis of CRF would help ameliorate the negative effects of CRF, which would mimic the regulatory role of cortisol.^{1,20} This is, perhaps, one of the most likely mechanisms of actions of herbs having adaptogenic properties. Thus, adaptogen remedies may be mimicking cortisol, adrenaline, or possibly ACTH or CRF, in a way that does not cause the same deleterious effects in stress dysfunction cascades, yet somehow tricks the body into thinking these endogenous substances are being supplied in a normal way. This mechanism may modulate allostasis and lower allostatic overload.^{25,31}

Remedies with an adaptogenic property must attenuate or prevent the initial dysfunction of neurotransmitters and hormones by increasing resistance to stress and in addition, attenuate or prevent pathological conditions resulting from the dysfunction.^{23,32} Adaptogens increase the organisms ability to maintain both homeostasis and allostasis.³³

In the present study we have evaluated the efficacy of *Ocimum sanctum* (leaves), *Withania somnifera* (roots), *Bacopa monnieri* (leaves) in the prevention of stress-induced changes during swim endurance test and cold restraint stress in mice. The standard group was administered water soluble panax root powder (*Panax ginseng*) orally while the control group was treated with distilled water alone. Stress is known to impair the metabolic functions of the body.^{30,34} The cold restraint stress-induced animals showed increased plasma cortisol levels. Such stress-induced biochemical impairment was attenuated by *Ocimum sanctum*, *Withania somnifera*, *Bacopa monnieri* extract treatment.^{12–15,17–20,25–28} In the swim endurance test also, increased swimming time was observed in mice pretreated with adaptogenic herbs. Mice pretreated with adaptogens showed significant improvement in the swimming time suggesting a central nervous system stimulant.^{33,35} Therefore, pretreatment of

mice with ethanolic extract of *Ocimum sanctum* 47 mg/kg, *Withania somnifera* 23 mg/kg, *Bacopa monnieri* 27 mg/kg p.o reduced the elevated plasma cortisol level as well as increased swim endurance in mice showing adaptogenic and anti-stress activity.^{30,36}

6. Conclusion

Cortisol is often referred to as the stress hormone and it is involved in the response to stress.²⁵ As cortisol has an immunosuppressive action, balance and systemic equilibrium of the adrenal glands can in turn, positively contribute to good metabolic rate, a strong immune system, healthy energy levels and normal sleep states.²⁰ Overall health, as well as the well being and optimum functioning of the adrenal glands should be addressed. Relaxation methods, as well as eating a healthy, balanced diet and exercising regularly can all be of great benefit to adrenal health and harmony.¹¹ Natural remedies have been used in traditional medicine for thousands of years to support the solid health of the adrenal glands, helping to encourage healthy and efficient adrenal function.^{1,4,5}

7. Source of Funding

None.

8. Conflict of Interest

The authors declare no conflict of interest.

9. Acknowledgement

The author is thankful to the Department of Biochemistry, Patna University, Patna, for all necessary support.

References

- Panosian A, Wikman G, Wagner H. Plant adaptogens, III, Earlier and more recent aspects and concepts on their mode of action. *Phytomedicine*. 1999;6(4):287–300.
- Selye H. Studies on Adaptation. *Endocrinology*. 1937;21(2):169–88.
- Selye H. Adaptation Energy. *Nature*. 1938;141:926. doi:10.1038/141926a0.
- Brekman II. Man and Biologically Active Substances. Pergamon; 1980.
- Selye H. A Syndrome produced by Diverse Noxious Agents. *Nature*. 1936;138:32. doi:10.1038/138032a0.
- Selye H. Experimental evidence supporting the conception of 'adaptation energy'. *Am J Physiol*. 1938;123:758–65.
- Bhattacharya SK, Goel RK, Kaur R, Ghosal S. Anti-stress activity of sitoindosides VII and VIII, new acylsterylglucosides from *Withania somnifera*. *Phytotherapy Res*. 1987;1(1):32–7. doi:10.1002/ptr.2650010108.
- Ghosal S, Lal J, Srivastava R, Bhattacharya SK, Upadhyay SN, Jaiswal AK, et al. Immunomodulatory and CNS effects of sitoindosides IX and X, two new glycowithanolides from *Withania somnifera*. *Phytotherapy Res*. 1989;3(5):201–6. doi:10.1002/ptr.2650030510.
- Godhwani S, Godhwani JL, Vyas DS. *Ocimum sanctum*: an experimental study evaluating its anti-inflammatory, analgesic and antipyretic activity in animals. *J Ethnopharmacol*. 1987;21(2):153–63. doi:10.1016/0378-8741(87)90125-5.
- Gupta S, Aslakson E, Gurbaxani BM. Inclusion of the glucocorticoid receptor in a hypothalamic pituitary adrenal axis model reveals bistability. *Theor Biol Med Model*. 2007;4:8.
- Bove, Mary. Adrenal Function, Stress and Botanical Medicine. In: Medicines from the Earth Proceedings. Black Mountain, NC; 2003.
- Gupta SK, Prakash J, Srivastava S. Validation of traditional claim of *tulsi*, *Ocimum sanctum* Linn. as a medicinal plant. *Indian J Exp Biol*. 2002;40:765–73.
- Bhargava KP, Singh N. Anti-stress activity of *Ocimum sanctum* Linn. *Indian J Med Res*. 1981;73:443–51.
- Devi U, Radioprotective P. anticarcinogenic and antioxidant properties of the Indian holy basil, *Ocimum sanctum* (Tulsi). *Indian J Exp Biol*. 2001;39:185–90.
- Satyavati GV, Gupta AK, Tandon N. *Ocimum sanctum* Linn. In: Medicinal Plants of India. India: ICMR; 1987. p. 355–71.
- Dhuley J. Adaptogenic and Cardioprotective action of ashwagandha in rats and frogs. *J Ethnopharmacol*. 2000;70(1):57–63. doi:10.1016/S0378-8741(99)00177-4.
- Archana R, Namasivayam A. Anti-stressor effect of *Withania somnifera*. *J Ethnopharmacol*. 1999;64:91–3. doi:10.1016/S0378-8741(98)00107-x.
- Bhattacharya SK, Goel RK, Kaur R, Ghosal S. Anti-stress activity of sitoindosides VII and VIII, new acylsterylglucosides from *Withania somnifera*. *Phytother Res*. 1987;1(1):32–7. doi:10.1002/ptr.2650010108.
- Bhattacharya SK, Muruganandam AV. Adaptogenic activity of *Withania somnifera*: an experimental study using a rat model of chronic stress. *Biochem Behav*. 2003;75:547–55.
- Biswas NM, Sengupta R, Roychoudhuri G, Chattopadhyay A, Sarkar M. Prevention of adrenocortical hyperactivity by dietary casein in rats exposed to forced swimming stress. *Indian J Exp Biol*. 2001;39:178–80.
- Nathan PJ, Clarke J, Lloyd J, Hutchison CW, Downey L, Stough C. The acute effects of an extract of *B. monnieri* (Brahmi) on cognitive function in healthy normal subjects. *Hum Psychopharmacol*. 2001;16(4):345–51.
- Stough C, Lloyd J, Downey LA, Hutchison CW, Rodgers T, Nathan PJ. The chronic effects of an extract of *B. monnieri* (Brahmi) on cognitive function in healthy normal subjects. *Psychopharmacology (Berl)*. 2001;156(4):481–4. doi:10.1007/s002130100815.
- Singh N, Misra N, Srivastava AK, Dixit KS, Gupta GP. Effects of anti-stress plants on biochemical changes during stress reaction. *Indian J of Pharmacol*. 1991;23:137–42.
- Singh RH, Singh L. Studies on the anti-anxiety effect of the Medhya Rasayana drug, Brahmi (*B. monnieri* Wettst.) - Part 1. *J Res Ayur Siddha*. 1980;1:133–48.
- George P, Chrousos MD, Philip W, Gold MD. The concept of stress system disorders. *J Am Med Assn*. 1992;267:1244–52.
- Rai D, Bhatia G, Palit G, Pal R, Singh S, Singh HK. Adaptogenic effect of *Bacopa monnieri* (Brahmi). *Pharmacol Biochem Behav*. 2003;75(4):823–30.
- Sairam K, Dorababu M, Goel RK, Bhattacharya SK. Antidepressant activity of standardized extract of *B. monnieri* in experimental models of depression in rats. *Phytomedicine*. 2002;9(3):207–11. doi:10.1078/0944-7113-00116.
- Sharma R, Chaturvedi C, Tiwari PV. Efficacy of *B. monnieri* in revitalizing intellectual functions in children. *J Res Edu Indian Med*. 1987;p. 1–12.
- Bone K. Clinical Applications of Ayurvedic and Chinese Herbs: Monographs for the Western Herbal Practitioner. Warwick, Queensland: Phytotherapy Press; 1996.
- Chrousos GP. Stressors, stress, and neuroendocrine integration of the adaptive response. The 1997 Hans Selye Memorial Lecture. *Ann N Y Acad Sci*. 1998;851:311–35.
- Wagner H, Norr H, Winterhoff H. Plant Adaptogens. *Phytomedicine*. 1994;1(1):63–76.
- Yance D. Adaptogens: New Conceptions and Userpersonal Insights, and Recent Advances, Centre for Natural Healing, 2000.

33. Carlini EA. Plants and the central nervous system. *Pharmacol Biochem Behav.* 2003;75:501–12.
34. Chrousos G, Gold PW. The concepts of stress and stress system disorders. *J Am Med Assoc.* 1992;267:1244–52.
35. Carrasco GA, Kar LD. Neuroendocrine Pharmacology of stress. *Eur J Pharmacol.* 2003;463:235–72. doi:10.1016/s0014-2999(03)01285-8.
36. Brekhman II, Dardymov IV. New substances of plant origin which increase non-specific resistance. *Ann Rev Pharmacol.* 1969;9:419–30.

Author biography

Anju, Assistant Professor

Ashis Kumar Ghosh, Former Professor and Head

Cite this article: Anju, Kumar Ghosh A. Adaptogens as anti-stress agents in reducing increased plasma cortisol level during stress. *Int J Clin Biochem Res* 2021;8(3):198-203.